


Math and Climate Seminar IMA




Mathematics and Climate Research Network


Joint MCRN/IMA Math and Climate Seminar
Tuesdays 11:15 – 12:05
streaming video available at
www.ima.umn.edu

MCRN www.mathclimate.org

An Introduction to Energy Balance Models
Richard McGehee



Seminar on the Mathematics of Climate
IMA, MCRN, School of Mathematics
October 1, 2013



Energy Balance


Conservation of Energy

temperature change \sim energy in – energy out

short wave energy from the Sun

long wave energy from the Earth

Everything else is detail.




Energy Balance

Stefan-Boltzmann Law

$$F = \sigma T^4$$

power flux (W/m²) temperature (K)

Stefan-Boltzmann constant
 $\sigma \approx 5.67 \times 10^{-8} \text{ W/m}^2 \text{ K}^4$



Energy Balance


Stefan-Boltzmann Law

$$F = \sigma T^4$$


power flux (W/m²) temperature (K)

Stefan-Boltzmann constant
 $\sigma \approx 5.67 \times 10^{-8} \text{ W/m}^2 \text{ K}^4$

Example
surface temperature of the Sun: 5780K
power flux: $5.67 \times 10^{-8} \times (5780)^4 = 6.33 \times 10^7 \text{ W/m}^2$
total solar power output: $6.33 \times 10^7 \times 4\pi(r_s)^2$,
where r_s = radius of the sun = $6.96 \times 10^8 \text{ m}$
total solar output: $3.85 \times 10^{26} \text{ W}$
200 nanoseconds = time it takes for the sun to produce the equivalent of the annual global electricity production



<http://astronomybythecosmos.com/tag/sun/>



Energy Balance

Insolation

Solar flux at a distance r from the sun:

$$F = \frac{6.33 \times 10^7 \cdot 4\pi r_s^2}{4\pi r^2} = 6.33 \times 10^7 \left(\frac{r_s}{r}\right)^2 \text{ W/m}^2$$

$r_s = 6.96 \times 10^8 \text{ m}$
 $r = 1.5 \times 10^{11} \text{ m}$
 $F = 1368 \text{ W/m}^2$

Power intercepted by the Earth:

$$F \times \pi r_e^2 \text{ W}, \quad r_e = \text{radius of Earth} = 6.37 \times 10^6 \text{ m}$$

$$F = 1.74 \times 10^{17} \text{ W}$$

Energy Balance
Insolation


Solar flux at a distance r from the sun:

$$F = \frac{6.33 \times 10^7 4\pi r_s^2}{4\pi r^2} = 6.33 \times 10^7 \left(\frac{r_s}{r}\right)^2 \text{ W/m}^2$$

$r_s = 6.96 \times 10^8 \text{ m}$
 $r = 1.5 \times 10^{11} \text{ m}$
 $F = 1368 \text{ W/m}^2$

Power intercepted by the Earth:
 $F \times \pi r_E^2 \text{ W}$, $r_E = \text{radius of Earth} = 6.37 \times 10^6 \text{ m}$
 $F = 1.74 \times 10^{17} \text{ W}$

Biologically Stored Energy
total coal reserves: 10^{15} kg
energy content: $3 \times 10^7 \text{ J/kg}$
total energy in coal reserves: $3 \times 10^{22} \text{ J}$
 $= 2 \text{ days of insolation}$



Energy Balance
Insolation

Global Average Insolation
intercepted flux: $F = 1368 \text{ W/m}^2$
Earth cross-section: πr_E^2
surface area: $4\pi r_E^2$
average flux: $1368/4 = 342 \text{ W/m}^2 = Q$

Simple Model
Assume that Earth is a perfectly thermally conducting black body.

$$Q = \sigma T^4$$

$$T = (Q/\sigma)^{1/4} = (342/5.67 \times 10^{-8})^{1/4}$$

$$= 279\text{K} = 6^\circ\text{C} = 43^\circ\text{F}$$

Dynamics
 $R \frac{dT}{dt} = Q - \sigma T^4$

heat capacity \rightarrow \leftarrow stable equilibrium

Energy Balance
Albedo

Not all the insolation reaches the surface. Some is reflected back into space.
The proportion reflected is called the albedo, denoted α .
For Earth, $\alpha \approx 0.3$.

Simple Model
Assume that Earth is a perfectly thermally conducting black body, but only 70% of the insolation is absorbed.

$$T = (0.7 \cdot F/\sigma)^{1/4} = (0.7 \cdot 342/5.67 \times 10^{-8})^{1/4}$$

$$= 255\text{K} = -18^\circ\text{C} = 0^\circ\text{F}$$

Dynamics
 $R \frac{dT}{dt} = Q(1 - \alpha) - \sigma T^4$ \leftarrow stable equilibrium


Energy Balance
Albedo

Not all the insolation reaches the surface. Some is reflected back into space.
The proportion reflected is called the albedo, denoted α .
For Earth, $\alpha \approx 0.3$.

Simple Model
Assume that Earth is a perfectly thermally conducting black body, but only 70% of the insolation is absorbed.


Why isn't the Earth a snowball?

$$T = (0.7 \cdot F/\sigma)^{1/4} = (0.7 \cdot 342/5.67 \times 10^{-8})^{1/4}$$

$$= 255\text{K} = -18^\circ\text{C} = 0^\circ\text{F}$$


Energy Balance
Black Body Radiation
Planck's Function


frequency \rightarrow Planck's constant
flux density $\rightarrow B(\nu, T) = \frac{2h\nu^3}{c^2} \frac{1}{e^{h\nu/kT} - 1}$
temperature \rightarrow speed of light Boltzmann's constant



Flux density ($\text{W m}^{-2} \mu\text{m}^{-1} \text{cm}^{-1}$) vs Wavenumber (cm^{-1})




Raymond T. Pierrehumbert, *Principles of Planetary Climate*, Cambridge University Press, 2010.

Energy Balance
Insolation vs. OLR
OLR = Outgoing Longwave Radiation



Normalized spectrum vs Wavenumber (cm^{-1})

Raymond T. Pierrehumbert, *Principles of Planetary Climate*, Cambridge University Press, 2010.



Energy Balance

OLR as a Function of Surface Temperature

$$OLR = A + BT$$

A and B are determined from satellite observations.
 T is surface temperature (in Celsius).

$$A = 202 \text{ W/m}^2$$

$$B = 1.90 \text{ W/m}^2\text{K}$$

Kelvin → Dynamics → $R \frac{dT}{dt} = Q(1 - \alpha) - \sigma T^4$ → photosphere temperature

Celsius → becomes → $R \frac{dT}{dt} = Q(1 - \alpha) - (A + BT)$ → global mean surface temperature

Energy Balance

Homogeneous Earth

$$R \frac{dT}{dt} = Q(1 - \alpha) - (A + BT)$$

Equilibrium Temperature

$$T_{eq} = \frac{Q(1 - \alpha) - A}{B}$$

Stable, since $B > 0$.

Ice-free planet: $\alpha = 0.32$, $T_{eq} = 16^\circ\text{C}$
 Snowball planet: $\alpha = 0.62$, $T_{eq} = -38^\circ\text{C}$
 No glacier would form on an ice-free Earth.
 No glacier would melt on a snowball Earth.

Easy question: **Why do we have ice caps?**
 Hard question: **If Earth was ever a snowball, how did we get out?**

Energy Balance

Latitude Dependence

Make T depend on $y = \sin(\text{latitude})$

$$R \frac{\partial T(y,t)}{\partial t} = Qs(y)(1 - \alpha) - (A + BT(y,t))$$

insolation distribution

Q = global annual average insolation = 342 W/m^2
 $s(y)$ = distribution across latitudes ($\int_0^1 s(y) dy = 1$)







One can show that


$$s(y) = \frac{2}{\pi} \int_0^{2\pi} \sqrt{1 - (1 - y^2) \sin^2 \beta \cos^2 \theta - y \cos \beta} d\theta$$

β = obliquity = 23.5°

Chylek and Coakley's quadratic approximation:

$$s(y) \approx 1 - 0.241(3y^2 - 1)$$



 **Energy Balance**


Budyko's Equation

$$R \frac{\partial T}{\partial t} = Qs(y)(1-\alpha) - (A+BT) + C(\bar{T}-T)$$


$$\bar{T}(t) = \int_0^1 T(y,t) dy$$

Weather

Second Law of Thermodynamics:
Energy travels from hot places to cold places.



Budyko's equation as a dynamical system:
 T lives in a function space (temperature as a function of latitude).

 **Energy Balance**

Budyko's Equilibrium

$$R \frac{\partial T}{\partial t} = Qs(y)(1-\alpha(y)) - (A+BT) + C(\bar{T}-T)$$

albedo depends on latitude

equilibrium solution: $T = T^*(y)$

$$Qs(y)(1-\alpha(y)) - (A+BT^*(y)) + C(\bar{T} - T^*(y)) = 0$$

Integrate:


$$\int_0^1 (Qs(y)(1-\alpha(y)) - (A+BT^*(y)) + C(\bar{T} - T^*(y))) dy = 0$$

$$Q(1-\bar{\alpha}) - (A+BT^*) = 0$$

where $\bar{\alpha} = \int_0^1 \alpha(y)s(y) dy$

Global mean temperature at equilibrium

$$\bar{T} = \frac{1}{B}(Q(1-\bar{\alpha}) - A)$$

 **Energy Balance**

Budyko's Equilibrium

$$Qs(y)(1-\alpha(y)) - (A+BT^*(y)) + C(\bar{T} - T^*(y)) = 0$$


Global mean temperature at equilibrium:


$$\bar{T} = \frac{1}{B}(Q(1-\bar{\alpha}) - A) \quad \left(\bar{\alpha} = \int_0^1 \alpha(y)s(y) dy \right)$$

Equilibrium temperature profile:

$$T^*(y) = \frac{1}{B+C} (Qs(y)(1-\alpha(y)) - A + C\bar{T})$$

$C = 3.04$
 $\alpha = 0.32$: ice free
 $\alpha = 0.62$: snowball



 **Energy Balance**

Budyko's Equation

$$R \frac{\partial T}{\partial t} = Qs(y)(1-\alpha(y)) - (A+BT) + C(\bar{T}-T)$$

Next Time:

Budyko's equation as a model of glacial cycles.